

A RIGIDITY RESULT FOR NONLOCAL SEMILINEAR EQUATIONS

ALBERTO FARINA AND ENRICO VALDINOCI

ABSTRACT. We consider a possibly anisotropic integro-differential semilinear equation, run by a nondecreasing nonlinearity. We prove that if the solution grows at infinity less than the order of the operator, then it must be affine (possibly constant).

1. INTRODUCTION

It dates back to Liouville and Cauchy in 1844 that bounded harmonic functions are constant. Several generalizations of this result appeared in the literature, also involving nonlinear equations and more general growth of the solution at infinity (see [F] for a detailed review of this topic).

The purpose of this note is to obtain a rigidity result for integro-differential semilinear equations of fractional order $2s$, with $s \in (0, 1)$.

We recall that fractional integro-differential operators are a classical topic in analysis, whose study arises in different fields, including harmonic analysis [St], partial differential equations [C] and probability [Be]. The study of these operators has also relevance in concrete situations, in view of the related real-world applications, such as quantum mechanics [FLl], water waves [CSS], meteorology [CV], crystallography [G], biology [AAVV], finance [Sc] and high technology [ZL], just to name a few.

The type of integro-differential operators that we consider here are of the form

$$(1) \quad \mathcal{I}u(x) := \int_{S^{n-1} \times \mathbb{R}} (u(x + \vartheta r) + u(x - \vartheta r) - 2u(x)) \frac{d\mu(\vartheta) dr}{|r|^{1+2s}}.$$

In this notation, $r \in \mathbb{R}$ gets integrated by the usual Lebesgue measure dr , while $\vartheta \in S^{n-1}$ is integrated by a measure $d\mu(\vartheta)$, which is called “spectral measure” in jargon, and that satisfy the following nondegeneracy assumptions: there exist $\lambda, \Lambda \in (0, +\infty)$ such that

$$(2) \quad \inf_{\nu \in S^{n-1}} \int_{S^{n-1}} |\nu \cdot \vartheta|^{2s} d\mu(\vartheta) \geq \lambda \quad \text{and} \quad \mu(S^{n-1}) \leq \Lambda.$$

2010 *Mathematics Subject Classification.* 35R11, 35B53, 35R09.

Key words and phrases. Nonlocal integro-differential semilinear equations, Liouville-type theorems, nondecreasing nonlinearities.

The authors have been supported by the ERC grant 277749 “EPSILON Elliptic Pde’s and Symmetry of Interfaces and Layers for Odd Nonlinearities”.

Particularly famous cases of spectral measures are the ones induced by singular kernels, i.e. when $d\mu(\vartheta) = \mathcal{K}_0(\vartheta) d\mathcal{H}^{n-1}(\vartheta)$, with $0 < \inf_{S^{n-1}} \mathcal{K}_0 \leq \sup_{S^{n-1}} \mathcal{K}_0 < +\infty$. Notice that in this particular case the spectral measure is absolutely continuous with respect to the standard Hausdorff measure on S^{n-1} , and the operator in (1) comes from the integration against the homogeneous kernel

$$(3) \quad \mathcal{K}(y) := |y|^{-n-2s} \mathcal{K}_0 \left(\frac{y}{|y|} \right),$$

in the sense that

$$(4) \quad \mathcal{I}u(x) = \int_{\mathbb{R}^n} (u(x+y) + u(x-y) - 2u(x)) \mathcal{K}(y) dy.$$

Of course, the case $\mathcal{K}$ equal to constant boils down to the fractional Laplacian, i.e. to the case in which, up to a normalization factor, $\mathcal{I} = -(-\Delta)^s$.

The literature has recently shown an increasing effort towards the study of these types of anisotropic operators, see e.g. [KRS, KS, ROS, ROV, ROSV]. It is worth recalling that the picture provided by the general operator in (1) is often quite special when compared with the isotropic case, and sometimes even surprising: for instance, a complete regularity theory in the setting of (1)–(2) does not hold, and explicit counterexamples can be constructed, see [ROV].

We will consider the equation $\mathcal{I}u = f(u)$. This type of equations is often called “semilinear” since the nonlinearity only depends on the values of the solution itself (for these reasons, solutions of semilinear equations may satisfy better geometric properties than solutions of arbitrary equations).

Our main result states that if f is nondecreasing, then solutions of $\mathcal{I}u = f(u)$ whose growth at infinity is bounded by $|x|^\kappa$, with κ less than the order of operator, must be necessarily affine (and, in fact, constant when the nonlinearity is nontrivial). More precisely, we have:

Theorem 1. *Let $f \in C(\mathbb{R})$ be nondecreasing. Let $u \in C^2(\mathbb{R}^N)$ be a solution of*

$$(5) \quad \mathcal{I}u(x) = f(u(x)) \quad \text{for any } x \in \mathbb{R}^n.$$

Assume that

$$(6) \quad |u(x)| \leq K(1 + |x|^\kappa),$$

for some $K \geq 0$ and $\kappa \in [0, 2s)$.

Then the following classification holds true:

- (i) *if f is not identically zero, then u is constant, say $u(x) = c$ for any $x \in \mathbb{R}^n$, and $f(c) = 0$;*
- (ii) *if f is identically zero, then u is an affine function, say $u(x) = \varpi \cdot x + c$, for some $\varpi \in \mathbb{R}^n$ and $c \in \mathbb{R}$; in this case, if additionally $\kappa < 1$, then $\varpi = 0$ and u is constant.*

We remark that Theorem 1 holds true here for the very general integro-differential operator in (1)–(2). Moreover, as far as we know, Theorem 1 is new even in the case of regular spectral measure in (3)–(4), and, perhaps quite surprisingly, even in the isotropic case of the fractional Laplacian.

On the other hand, when $\mathcal{I}$ is replaced by the Laplacian (which is formally the above case with $s = 1$) Theorem 1 is a well known result in the framework of classical Liouville-type theorems: see for instance [F, Se]. As a matter of fact, the counterpart of (5) in the classical case, is the semilinear equation $\Delta u = f(u)$, set in the whole of $\mathbb{R}^n$. This equation has been extensively studied, also in connection with phase transition models, see e.g. [M, Br] and references therein. Its fractional analogue has also physical relevance, since it appears, for instance, in the study of phase transitions arising from long-range interactions and in the dynamics of atom dislocations in crystal, see e.g. [CSM, SV, CSir, GM, MP, DPV].

The strategy of the proof of Theorem 1 is to show that $f(u(x))$ must be identically equal to zero in any case, therefore u is a solution of $\mathcal{I}u = 0$, and this will allow us to use a Liouville-type theorem in order to obtain the desired classification. To this goal, one uses the subcritical growth of the solution u to compare with suitable barriers. Of course, the construction of the appropriate barriers is the main novelty with respect to the classical case, since the nonlocality of the operator mostly comes into play in this framework.

It is also worth pointing out that Theorem 1 has a natural, and simple, generalization that deals with the case in which (6) is replaced by a one-side inequality. In this spirit, we present the following result:

Theorem 2. *Let $f \in C(\mathbb{R})$ be nondecreasing. Let $u \in C^2(\mathbb{R}^N)$ be a solution of*

$$\mathcal{I}u(x) = f(u(x)) \quad \text{for any } x \in \mathbb{R}^n.$$

Then, if

$$u(x) \leq K(1 + |x|^\kappa),$$

for some $K \geq 0$ and $\kappa \in [0, 2s)$, we have that

$$\mathcal{I}u(x) \leq 0 \quad \text{for any } x \in \mathbb{R}^n.$$

Similarly, if

$$u(x) \geq -K(1 + |x|^\kappa),$$

for some $K \geq 0$ and $\kappa \in [0, 2s)$, we have that

$$\mathcal{I}u(x) \geq 0 \quad \text{for any } x \in \mathbb{R}^n.$$

We also point out that the regularity assumptions of u in Theorems 1 and 2 were taken for the sake of simplicity and, in concrete cases, one does not need u to be smooth to start with (for instance, in the setting of [KRS] one can deal with viscosity solutions and in the setting of [ROS] one can deal with weak solutions).

The rest of the paper is organized as follows. First, in Section 2 we collect some preliminary integral computations that will be used in Section 3 to construct a useful barrier. Roughly speaking, this barrier replaces the classical paraboloid in our nonlocal framework (of course, checking the properties of the paraboloid in the classical case is much simpler than constructing barriers in nonlocal cases).

The proofs of Theorems 1 and 2 occupy Section 4.

2. TOOLBOX

Below are some preliminary integral computations, needed to construct a suitable barrier in Section 3. For convenience, we use the notation

$$\mathcal{I}_1 v(x) := \int_{S^{n-1} \times (-1,1)} (v(x + \vartheta r) + v(x - \vartheta r) - 2v(x)) \frac{d\mu(\vartheta) dr}{|r|^{1+2s}}$$

$$\text{and } \mathcal{I}_2 v(x) := \int_{S^{n-1} \times (\mathbb{R} \setminus (-1,1))} (v(x + \vartheta r) + v(x - \vartheta r) - 2v(x)) \frac{d\mu(\vartheta) dr}{|r|^{1+2s}},$$

in order to distinguish the integration performed when $|r| < 1$ from the one when $|r| \geq 1$.

2.1. Estimates near the origin. Here we estimate $\mathcal{I}_1 v$ and $\mathcal{I}_2 v$ near the origin according to the following Lemmata 1 and 2:

Lemma 1. *Let $v \in C^2(B_3)$. Then, for any $x \in B_1$,*

$$\mathcal{I}_1 v(x) \leq C,$$

for some $C > 0$ possibly depending on n, s, Λ and $\|v\|_{C^2(B_2)}$.

Proof. If $x, y \in B_1$ we obtain from a Taylor expansion that

$$|v(x + y) + v(x - y) - 2v(x)| \leq \|D^2 v\|_{L^\infty(B_2)} |y|^2.$$

hence, by integration, we get

$$\begin{aligned} \mathcal{I}_1 v(x) &\leq \int_{S^{n-1} \times (-1,1)} \|D^2 v\|_{L^\infty(B_2)} |r|^2 \frac{d\mu(\vartheta) dr}{|r|^{1+2s}} \\ &\leq \Lambda \|D^2 v\|_{L^\infty(B_2)} \int_0^1 |r|^{1-2s} dr, \end{aligned}$$

thanks to (2), which gives the desired result. $\square$

Lemma 2. *Let*

$$(7) \quad \gamma \in (0, 2s).$$

Let $v : \mathbb{R}^n \rightarrow [0, +\infty)$ be a measurable function such that $v(x) \leq |x|^\gamma$ for any $x \in \mathbb{R}^n$. Then, for any $x \in B_1$,

$$\mathcal{I}_2 v(x) \leq C,$$

for some $C > 0$ possibly depending on n, s, Λ and γ .

Proof. Let $x \in B_1$ and $y \in \mathbb{R}^n \setminus B_1$. Then $|x| \leq 1 \leq |y|$ and so

$$\begin{aligned} & |v(x+y) + v(x-y) - v(x)| \\ & \leq |v(x+y)| + |v(x-y)| + |v(x)| \\ & \leq |x+y|^\gamma + |x-y|^\gamma + |x|^\gamma \\ & \leq 2(|x| + |y|)^\gamma + |x|^\gamma \\ & \leq (2^{\gamma+1} + 1) |y|^\gamma. \end{aligned}$$

So, we integrate, we recall (2) and we see that

$$\begin{aligned} \mathcal{I}_2 v(x) & \leq \int_{S^{n-1} \times (\mathbb{R}^n \setminus (-1,1))} (2^{\gamma+1} + 1) |r|^\gamma \frac{d\mu(\vartheta) dr}{|r|^{1+2s}} \\ & \leq 2(2^{\gamma+1} + 1) \Lambda \int_1^{+\infty} r^{\gamma-1-2s} dr. \end{aligned}$$

Then we use (7) and we obtain the desired result. $\square$

2.2. Estimates far from the origin. Now we estimate $\mathcal{I}v = \mathcal{I}_1 v + \mathcal{I}_2 v$ at infinity:

Lemma 3. *Let γ be as in (7) and $v : \mathbb{R}^n \rightarrow \mathbb{R}$ be a measurable function such that $v(x) \leq |x|^\gamma$ for any $x \in \mathbb{R}^n$.*

Assume also that $v(x) = |x|^\gamma$ for any $x \in \mathbb{R}^n \setminus B_1$. Then, for any $x \in \mathbb{R}^n \setminus B_1$,

$$\mathcal{I}v(x) \leq C,$$

for some $C > 0$ possibly depending on n, s, Λ and γ .

Proof. Fix $x \in \mathbb{R}^n \setminus B_1$. Then $v(x) = |x|^\gamma$. Moreover $v(x \pm y) \leq |x \pm y|^\gamma$, and so

$$v(x+y) + v(x-y) - 2v(x) \leq |x+y|^\gamma + |x-y|^\gamma - 2|x|^\gamma.$$

Therefore, calling $\omega := x/|x|$ and changing variable $r := |x|\varrho$, we have that

$$\begin{aligned} \mathcal{I}v(x) & = \int_{S^{n-1} \times \mathbb{R}} (v(x + \vartheta r) + v(x - \vartheta r) - 2v(x)) \frac{d\mu(\vartheta) dr}{|r|^{1+2s}} \\ & \leq \int_{S^{n-1} \times \mathbb{R}} (|x + \vartheta r|^\gamma + |x - \vartheta r|^\gamma - 2|x|^\gamma) \frac{d\mu(\vartheta) dr}{|r|^{1+2s}} \\ (8) \quad & = |x|^{\gamma-2s} \int_{S^{n-1} \times \mathbb{R}} (|\omega + \vartheta \varrho|^\gamma + |\omega - \vartheta \varrho|^\gamma - 2) \frac{d\mu(\vartheta) d\varrho}{|\varrho|^{1+2s}} \\ & = |x|^{\gamma-2s} \int_{S^{n-1} \times \mathbb{R}} \frac{g(\vartheta \varrho) + g(-\vartheta \varrho) - 2g(0)}{|\varrho|^{1+2s}} d\mu(\vartheta) d\varrho, \end{aligned}$$

where, for any $\eta \in \mathbb{R}^n$, we set $g(\eta) := |\omega + \eta|^\gamma$. Notice that

$$(9) \quad |g(\eta)| \leq (|\omega| + |\eta|)^\gamma = (1 + |\eta|)^\gamma.$$

Moreover $g \in C^\infty(B_{1/2})$ and, for any $\eta \in B_{1/2}$ we have that

$$\partial_i g(\eta) = \gamma |\omega + \eta|^{\gamma-2} (\omega_i + \eta_i)$$

$$\text{and} \quad \partial_{ij}^2 g(\eta) = \gamma(\gamma-2) |\omega + \eta|^{\gamma-4} (\omega_i + \eta_i)(\omega_j + \eta_j) + \gamma |\omega + \eta|^{\gamma-2} \delta_{ij}.$$

Consequently, for any $\eta \in B_{1/2}$,

$$|D^2 g(\eta)| \leq C_o |\omega + \eta|^{\gamma-2},$$

for some $C_o > 0$ depending on γ and n , and $|\omega + \eta| \geq |\omega| - |\eta| \geq 1/2$, therefore

$$\|D^2 g\|_{L^\infty(B_{1/2})} \leq 2^{2-\gamma} C_o.$$

This, together with a Taylor expansion, implies that, for any $\eta \in B_{1/2}$,

$$|g(\eta) + g(-\eta) - 2g(0)| \leq \|D^2 g\|_{L^\infty(B_{1/2})} |\eta|^2 \leq 2^{2-\gamma} C_o |\eta|^2.$$

Hence, recalling (9) and (2), we obtain that

$$\begin{aligned} & \int_{S^{n-1} \times \mathbb{R}} \frac{g(\vartheta \varrho) + g(-\vartheta \varrho) - 2g(0)}{|\varrho|^{1+2s}} d\mu(\vartheta) d\varrho \\ & \leq C \left(\int_{S^{n-1} \times (-1/2, 1/2)} \frac{|\varrho|^2}{|\varrho|^{1+2s}} d\mu(\vartheta) d\varrho + \int_{S^{n-1} \times (\mathbb{R} \setminus (-1/2, 1/2))} \frac{(1 + |\varrho|)^\gamma}{|\varrho|^{1+2s}} d\mu(\vartheta) d\varrho \right) \\ & \leq C\Lambda \left(\int_{(-1/2, 1/2)} \frac{|\varrho|^2}{|\varrho|^{1+2s}} d\varrho + \int_{\mathbb{R} \setminus (-1/2, 1/2)} \frac{(1 + |\varrho|)^\gamma}{|\varrho|^{1+2s}} d\varrho \right) \\ & \leq C'\Lambda, \end{aligned}$$

for some $C, C' > 0$, thanks to (7). We insert this into (8) and we obtain the desired estimate (by possibly renaming the constants). $\square$

3. CONSTRUCTION OF AN AUXILIARY BARRIER

Here we use the estimate in Section 2 and we borrow some ideas from [DSV] to construct a useful auxiliary function:

Lemma 4. *Let $\gamma \in (0, 2s)$. There exists a function $v \in C^\infty(\mathbb{R}^n)$ such that*

$$(10) \quad v(0) = 0,$$

$$(11) \quad 0 \leq v(x) \leq |x|^\gamma \text{ for any } x \in \mathbb{R}^n,$$

$$(12) \quad v(x) = |x|^\gamma \text{ if } |x| \geq 1$$

$$(13) \quad \text{and} \quad \sup_{x \in \mathbb{R}^n} \mathcal{I}v(x) \leq C,$$

for some $C > 0$.

Proof. Let $\tau \in C^\infty(\mathbb{R}^n)$ be such that $0 \leq \tau \leq 1$ in the whole of $\mathbb{R}^n$, $\tau = 1$ in $B_{1/2}$ and $\tau = 0$ in $\mathbb{R}^n \setminus B_1$. We define $v(x) := (1 - \tau(x))|x|^\gamma$. In this way, conditions (10), (11) and (12) are fulfilled.

Furthermore, v satisfies all the assumptions of Lemmata 1, 2 and 3. Thus, using such results, we obtain condition (13). $\square$

4. PROOF OF THE MAIN RESULTS

Proof of Theorem 1. The proof relies on a modification of a classical argument (see for instance [Se, F]). In our setting, the barrier constructed in Lemma 4 will replace (at least from one side) the classical paraboloid. The details of the argument goes as follows. Let f , u , K and κ as in the statement of Theorem 1. Let $\gamma := (2s + \kappa)/2$. By construction,

$$(14) \quad \gamma \in (\kappa, 2s),$$

so we can use the barrier v constructed in Lemma 4. We fix $\epsilon > 0$ and an arbitrary point $x_0 \in \mathbb{R}^n$, and we define

$$(15) \quad \begin{aligned} w_1(x) &:= u(x) - u(x_0) + 2\epsilon - \epsilon v(x - x_0) \\ \text{and} \quad w_2(x) &:= u(x) - u(x_0) - 2\epsilon + \epsilon v(x - x_0) \end{aligned}$$

We remark that

$$\begin{aligned} \limsup_{|x| \rightarrow +\infty} w_1(x) &\leq \limsup_{|x| \rightarrow +\infty} [u(x) + |u(x_0)| + 2\epsilon - \epsilon v(x - x_0)] \\ &\leq \limsup_{|x| \rightarrow +\infty} [K(1 + |x|^\kappa) + |u(x_0)| + 2\epsilon - \epsilon |x - x_0|^\gamma] = -\infty \\ \text{and} \quad \liminf_{|x| \rightarrow +\infty} w_2(x) &\geq \liminf_{|x| \rightarrow +\infty} [u(x) - |u(x_0)| - 2\epsilon + \epsilon v(x - x_0)] \\ &\geq \liminf_{|x| \rightarrow +\infty} [-K(1 + |x|^\kappa) - |u(x_0)| - 2\epsilon + \epsilon |x - x_0|^\gamma] = +\infty, \end{aligned}$$

where we have used (6), (12) and (14). As a consequence the maximum of w_1 and the minimum of w_2 are attained, i.e. there exists $y_1, y_2 \in \mathbb{R}^n$ such that

$$(16) \quad w_1(y) \leq w_1(y_1) \quad \text{and} \quad w_2(y) \geq w_2(y_2) \quad \text{for any } y \in \mathbb{R}^n.$$

Accordingly, for any $y \in \mathbb{R}^n$,

$$(17) \quad \begin{aligned} w_1(y_1 + y) + w_1(y_1 - y) - 2w_1(y_1) &\leq 0 \\ \text{and } w_2(y_1 + y) + w_2(y_1 - y) - 2w_2(y_2) &\geq 0. \end{aligned}$$

On the other hand

$$(18) \quad \begin{aligned} &w_1(y_1 + y) + w_1(y_1 - y) - 2w_1(y_1) \\ &= u(y_1 + y) + u(y_1 - y) - 2u(y_1) \\ &\quad - \epsilon(v(y_1 + y - x_0) + v(y_1 - y - x_0) - 2v(y_1 - x_0)), \\ \text{and} \quad &w_2(y_2 + y) + w_2(y_2 - y) - 2w_2(y_2) \\ &= u(y_2 + y) + u(y_2 - y) - 2u(y_2) \\ &\quad + \epsilon(v(y_2 + y - x_0) + v(y_2 - y - x_0) - 2v(y_2 - x_0)). \end{aligned}$$

By comparing (17) and (18), we obtain that

(19)

$$\begin{aligned} 0 &\geq \int_{S^{n-1} \times \mathbb{R}} (w_1(y_1 + \vartheta r) + w_1(y_1 - \vartheta r) - 2w_1(y_1)) \frac{d\mu(\vartheta) dr}{|r|^{1+2s}} \\ &= \mathcal{I}u(y_1) - \epsilon \mathcal{I}v(y_1 - x_0) \end{aligned}$$

and

$$\begin{aligned} 0 &\leq \int_{S^{n-1} \times \mathbb{R}} (w_2(y_2 + \vartheta r) + w_2(y_2 - \vartheta r) - 2w_2(y_2)) \frac{d\mu(\vartheta) dr}{|r|^{1+2s}} \\ &= \mathcal{I}u(y_2) + \epsilon \mathcal{I}v(y_2 - x_0). \end{aligned}$$

Therefore, using (5) and (13), we obtain that

$$(20) \quad 0 \geq f(u(y_1)) - C\epsilon \quad \text{and} \quad 0 \leq f(u(y_2)) + C\epsilon.$$

Now we observe that $w_1(x_0) = 2\epsilon \geq 0$ and $w_2(x_0) = -2\epsilon \leq 0$, thanks to (15) and (10). So, if we evaluate (16) at the point $y := x_0$, we obtain that

$$(21) \quad 0 \leq w_1(x_0) \leq w_1(y_1) \quad \text{and} \quad 0 \geq w_2(x_0) \geq w_2(y_2).$$

Furthermore, using that $v \geq 0$ (recall (11)), we see from (15) that

$$w_1(y_1) \leq u(y_1) - u(x_0) + 2\epsilon \quad \text{and} \quad w_2(y_2) \geq u(y_2) - u(x_0) - 2\epsilon.$$

By comparing this with (21), we conclude that

$$u(y_1) \geq u(x_0) - 2\epsilon \quad \text{and} \quad u(y_2) \leq u(x_0) + 2\epsilon.$$

Therefore, since f is nondecreasing, we deduce that

$$f(u(y_1)) \geq f(u(x_0) - 2\epsilon) \quad \text{and} \quad f(u(y_2)) \leq f(u(x_0) + 2\epsilon).$$

We plug this information into (20), and we obtain that

$$(22) \quad 0 \geq f(u(x_0) - 2\epsilon) - C\epsilon \quad \text{and} \quad 0 \leq f(u(x_0) + 2\epsilon) + C\epsilon.$$

We remark that x_0 was fixed at the beginning and so it is independent of ϵ (conversely, the points y_1 and y_2 in general may depend on ϵ). This says that we can pass to the limit as $\epsilon \rightarrow 0^+$ in (22) and use the continuity of f to obtain that

$$0 \geq f(u(x_0)) \quad \text{and} \quad 0 \leq f(u(x_0)),$$

that is $f(u(x_0)) = 0$. Since x_0 is an arbitrary point of $\mathbb{R}^n$, we have proved that

$$(23) \quad f(u(x)) = 0 \quad \text{for any } x \in \mathbb{R}^n.$$

Thus, using again (5), we obtain that

$$\mathcal{I}u = 0 \quad \text{in } \mathbb{R}^n.$$

From this and Theorem 2.1 in [ROS], we obtain that u is a polynomial of degree $d \in \mathbb{N}$, with d less than or equal to the integer part of κ . In particular, $d \leq \kappa < 2s < 2$, hence $d \in \{0, 1\}$ and thus u is an affine function. So we can write $u(x) = \varpi \cdot x + c$, for some $\varpi \in \mathbb{R}^n$ and $c \in \mathbb{R}$.

As a side remark notice that, since $d \leq \kappa$, when the additional assumption $\kappa < 1$ holds true, we have that $d = 0$, and consequently $\varpi = 0$ and u is constant. These considerations establish claim (ii) in Theorem 1.

Now we prove that

(24) if f does not vanish identically, then $\varpi = 0$.

Indeed, if, by contradiction, we had $\varpi \neq 0$, given any $r \in \mathbb{R}$, we can take $x_\star := (r - c)|\varpi|^{-2}\varpi$. Then

$$u(x_\star) = \varpi \cdot x_\star + c = r,$$

thus, by (23), we get that $f(r) = f(u(x_\star)) = 0$. Since r was arbitrary, this would say that f vanishes identically, in contradiction with our assumptions. This proves (24).

By (24) and (23) we obtain claim (i) in Theorem 1. This completes the proof of Theorem 1. $\square$

Proof of Theorem 2. The proof of Theorem 1 goes through in this case, just considering only the function w_1 (to obtain the first statement of Theorem 2), or only the function w_2 (to obtain the second statement). $\square$

REFERENCES

- [Be] JEAN BERTOIN, *Lévy processes*. Cambridge Tracts in Mathematics, Vol. 121, Cambridge University Press, Cambridge, 1996.
- [Br] ANDREA BRAIDES, *Γ -convergence for beginners*. Oxford Lecture Series in Mathematics and its Applications, Vol. 22, Oxford University Press, Oxford, 2002.
- [CSir] XAVIER CABRÉ AND YANNICK SIRE, *Nonlinear equations for fractional Laplacians. I: Regularity, maximum principles, and Hamiltonian estimates*. Ann. Inst. Henri Poincaré, Anal. Non Linéaire **31** (2014), no. 1, 23–53.
- [CSM] XAVIER CABRÉ AND JOAN SOLÀ-MORALES, *Layer solutions in a half-space for boundary reactions*. Commun. Pure Appl. Math. **58** (2005), no. 12, 1678–1732.
- [C] LUIS CAFFARELLI, *Further regularity for the Signorini problem*. Comm. Partial Differential Equations **4** (1979), no. 9, 1067–1075.
- [CV] LUIS CAFFARELLI AND ALEXIS VASSEUR, *Drift diffusion equations with fractional diffusion and the quasi-geostrophic equation*. Ann. Math. **171** (2010), no. 3, 1903–1930.
- [CSS] WALTER CRAIG, ULRICH SCHANZ AND CATHERINE SULEM, *The modulational regime of three-dimensional water waves and the Davey-Stewartson system*. Ann. Inst. H. Poincaré Anal. Non Linéaire **14** (1997), no 5, 615–667.
- [DPV] SERENA DIPIERRO, GIAMPIERO PALATUCCI, AND ENRICO VALDINOCI, *Dislocation dynamics in crystals: a macroscopic theory in a fractional Laplace setting*. Commun. Math. Phys. **333** (2015), no. 2, 1061–1105.
- [DSV] SERENA DIPIERRO, OVIDIU SAVIN AND ENRICO VALDINOCI, *A nonlocal free boundary problem*, preprint <http://arxiv.org/pdf/1411.7971.pdf> (2014).
- [F] ALBERTO FARINA, *Liouville-type theorems for elliptic problems*. Handbook of differential equations: stationary partial differential equations Vol. IV, 61–116, Elsevier/North-Holland, Amsterdam, 2007.
- [FLI] CHARLES FEFFERMAN AND RAFAEL DE LA LLAVE, *Relativistic stability of matter. I*. Rev. Mat. Iberoamericana **2** (1986), no. 1-2, 119–213.

- [G] GANG LU, *The Peierls-Nabarro model of dislocations: a venerable theory and its current development*. Handbook of materials modeling, 793–811, Springer, Sidney 2005.
- [GM] MARÍA DEL MAR GONZÁLEZ AND RÉGIS MONNEAU, *Slow motion of particle systems as a limit of a reaction-diffusion equation with half-Laplacian in dimension one*. Discrete Contin. Dyn. Syst. **32** (2012), no. 4, 1255–1286.
- [AAVV] NICOLAS E. HUMPHRIES ET AL., *Environmental context explains Lévy and Brownian movement patterns of marine predators*. Nature **465** (2010), no. 24, 1066–1069.
- [KRS] MORITZ KASSMANN, MARCUS RANG AND RUSSELL W. SCHWAB, *Integro-differential equations with nonlinear directional dependence*. Indiana Univ. Math. J. **63** (2014), no. 5, 1467–1498.
- [KS] MORITZ KASSMANN AND RUSSELL W. SCHWAB, *Regularity results for nonlocal parabolic equations*, preprint <http://arxiv.org/pdf/1305.5418.pdf> (2013).
- [M] Luciano Modica, *The gradient theory of phase transitions and the minimal interface criterion*. Arch. Ration. Mech. Anal. **98** (1987), 123–142.
- [MP] RÉGIS MONNEAU AND STEFANIA PATRIZI, *Homogenization of the Peierls-Nabarro model for dislocation dynamics*. J. Differ. Equations **253** (2012), no. 7, 2064–2105.
- [ROS] XAVIER ROS-OTON AND JOAQUIM SERRA, *Regularity theory for general stable operators*, preprint <http://arxiv.org/pdf/1412.3892.pdf> (2014).
- [ROSV] XAVIER ROS-OTON, JOAQUIM SERRA AND ENRICO VALDINOCI, *Pohozaev identities for anisotropic integro-differential operators*, preprint <http://arxiv.org/pdf/1502.01431.pdf> (2015).
- [ROV] XAVIER ROS-OTON AND ENRICO VALDINOCI, *The Dirichlet problem for non-local operators with singular kernels: convex and nonconvex domains*, preprint <http://arxiv.org/pdf/1502.00782.pdf> (2015).
- [Sc] WIM SCHOUTENS, *Lévy Processes in Finance. Pricing Financial Derivatives*. Wiley Series in Probability and Statistics, Wiley, England, 2003.
- [Se] JAMES SERRIN, *Entire solutions of nonlinear Poisson equations*. Proc. London. Math. Soc. (3) **24** (1972), 348–366.
- [SV] YANNICK SIRE AND ENRICO VALDINOCI, *Fractional Laplacian phase transitions and boundary reactions: a geometric inequality and a symmetry result*. J. Funct. Anal. **256** (2009), no. 6, 1842–1864.
- [St] ELIAS M. STEIN, *Singular integrals and differentiability properties of functions*. Princeton Mathematical Series, No. 30, Princeton University Press, Princeton N.J., 1970.
- [ZL] GANG ZHANG AND BAOWEN LI, *Anomalous vibrational energy diffusion in carbon nanotubes*. Journ. Chemical Phys. **123** (2005), no. 014705, 1–3.

ALBERTO FARINA: LAMFA – CNRS UMR 6140 – UNIVERSITÉ DE PICARDIE JULES VERNE – FACULTÉ DES SCIENCES – 33, RUE SAINT-LEU – 80039 AMIENS CEDEX 1, FRANCE – EMAIL: alberto.farina@u-picardie.fr

ENRICO VALDINOCI: WEIERSTRASS-INSTITUT FÜR ANGEWANDTE ANALYSIS UND STOCHASTIK – MOHRENSTRASSE, 39 – 10117 BERLIN, GERMANY – AND – UNIVERSITÀ DEGLI STUDI DI MILANO – DIPARTIMENTO DI MATEMATICA – VIA CESARE SALDINI, 50 – 20133 MILANO, ITALY – EMAIL: enrico@math.utexas.edu